

## Physics Formulas

---

Mechanics

Friction

Moment of Inertia

Newtonian Gravity

Projectile Motion

Simple Pendulum

Electricity

Thermodynamics

Electromagnetism

Optics

Quantum Physics

---

# Mechanics

---

Mechanics is the oldest branch of physics. Mechanics deals with all kinds and complexities of motion. It includes various techniques, which can simplify the solution of a mechanical problem.

## Motion in One Dimension

---

The formulas for motion in one dimension (Also called Kinematical equations of motion) are as follows. (Here 'u' is initial velocity, 'v' is final velocity, 'a' is acceleration and t is time):

Important topics of physics with equations

$$s = ut + \frac{1}{2} at^2$$

$$v = u + at$$

$$v^2 = u^2 + 2as$$

$$v_{av} \text{ (Average Velocity)} = \frac{(v+u)}{2}$$

## Momentum, Force and Impulse

---

Formulas for momentum, impulse and force concerning a particle moving in 3 dimensions are as follows (Here force, momentum and velocity are vectors ):

Momentum is the product of mass and velocity of a body. Momentum is calculate using the formula:  $P = m \text{ (mass)} \times v \text{ (velocity)}$

Force can defined as something which causes a change in momentum of a body. Force is given by the celebrated newton's law of motion:  $F = m \text{ (mass)} \times a \text{ (acceleration)}$

Impulse is a large force applied in a very short time period. The strike of a hammer is an impulse. Impulse is given by  $I = m(v-u)$

## Pressure

---

Pressure is defined as force per unit area:

$$\text{Pressure (P)} = \text{Force (F)}$$

$$\text{Area (A)}$$

## Density

---

Density is the mass contained in a body per unit volume.

The formula for density is:

$$\text{Density (D)} = \text{Mass(M)}$$

Volume (V)

## Angular Momentum

---

Angular momentum is an analogous quantity to linear momentum in which the body is undergoing rotational motion. The formula for angular momentum (J) is given by:

$$J = r \times p$$

where J denotes angular momentum, r is radius vector and p is linear momentum.

## Torque

---

Torque can be defined as moment of force. Torque causes rotational motion. The formula for torque is:  $\tau = r \times F$ , where  $\tau$  is torque, r is the radius vector and F is linear force.

## Circular Motion

---

The formulas for circular motion of an object of mass 'm' moving in a circle of radius 'r' at a tangential velocity 'v' are as follows:

$$\text{Centripetal force (F)} = mv^2$$

r

$$\text{Centripetal Acceleration (a)} = \frac{v^2}{r}$$

r

Center of Mass

General Formula for Center of mass of a rigid body is :

## Important topics of physics with equations

$$R = \frac{\sum Ni = 1}{m_i} r_i$$

$$\sum Ni = 1m_i$$

where R is the position vector for center of mass, r is the generic position vector for all the particles of the object and N is the total number of particles.

## Reduced Mass for two Interacting Bodies

The physics formula for reduced mass ( $\mu$ ) is :

$$\mu = \frac{m_1 m_2}{m_1 + m_2}$$

$$m_1 + m_2$$

where  $m_1$  is mass of the first body,  $m_2$  is the mass of the second body.

## Work and Energy

Formulas for work and energy in case of one dimensional motion are as follows:

$$W \text{ (Work Done)} = F \text{ (Force)} \times D \text{ (Displacement)}$$

Energy can be broadly classified into two types, Potential Energy and Kinetic Energy. In case of gravitational force, the potential energy is given by

$$P.E. \text{ (Gravitational)} = m \text{ (Mass)} \times g \text{ (Acceleration due to Gravity)} \times h \text{ (Height)}$$

The transitional kinetic energy is given by  $\frac{1}{2} m \text{ (mass)} \times v^2 \text{ (velocity squared)}$

## Important topics of physics with equations

### Power

Power is, work done per unit time. The formula for power is given as

$$\text{Power (P)} = \frac{W}{t}$$

$$P = I^2 R$$

where P=power, W = Work, t = time.

[[Back to Formula Index](#)]

# Friction

---

Friction can be classified to be of two kinds : Static friction and dynamic friction.

**Static Friction:** Static friction is characterized by a coefficient of static friction  $\mu_s$ . Coefficient of static friction is defined as the ratio of applied tangential force (F) which can induce sliding, to the normal force between surfaces in contact with each other. The formula to calculate this static coefficient is as follows:

$$\mu_s = \frac{\text{Applied Tangential Force (F)}}{\text{Normal Force(N)}}$$

Normal Force(N)

The amount of force required to slide a solid resting on flat surface depends on the coefficient of static friction and is given by the formula:

$$F_{\text{Horizontal}} = \mu_s \times M(\text{Mass of solid}) \times g \text{ (acceleration)}$$

Important topics of physics with equations

Dynamic Friction:

Dynamic friction is also characterized by the same coefficient of friction as static friction and therefore formula for calculating coefficient of dynamic friction is also the same as above. Only the dynamic friction coefficient is generally lower than the static one as the applied force required to overcome normal force is lesser.

[Back to Formula Index]

## Moment of Inertia

---

Here are some formulas for Moments of Inertia of different objects. (M stands for mass, R for radius and L for length):

Object Axis Moment of Inertia

Disk Axis parallel to disc, passing through the center  $MR^2/2$

Disk Axis passing through the center and perpendicular to disc  $MR^2/2$

Thin Rod Axis perpendicular to the Rod and passing through center  $ML^2/12$

Solid Sphere Axis passing through the center  $2MR^2/5$

Solid Shell Axis passing through the center  $2MR^2/3$

## Newtonian Gravity

---

Here are some important formulas, related to Newtonian Gravity:

Newton's Law of universal Gravitation:

Important topics of physics with equations

$$F_g = G \frac{m_1 m_2}{r^2}$$

$r^2$

where

$m_1, m_2$  are the masses of two bodies

$G$  is the universal gravitational constant which has a value of  $6.67300 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$

$r$  is distance between the two bodies

Formula for escape velocity ( $v_{esc}$ ) =  $(2GM / R)^{1/2}$  where,

$M$  is mass of central gravitating body

$R$  is radius of the central body

## Projectile Motion

---

Here are two important formulas related to projectile motion:

( $v$  = velocity of particle,  $v_0$  = initial velocity,  $g$  is acceleration due to gravity,  $\theta$  is angle of projection,  $h$  is maximum height and  $l$  is the range of the projectile.)

Maximum height of projectile ( $h$ ) =  $\frac{v_0^2 \sin^2 \theta}{2g}$

$2g$

Horizontal range of projectile ( $l$ ) =  $\frac{v_0^2 \sin 2\theta}{g}$

### Simple Pendulum

---

The physics formula for the period of a simple pendulum ( $T$ ) =  $2\pi \sqrt{l/g}$  where

$l$  is the length of the pendulum

Important topics of physics with equations

g is acceleration due to gravity

Conical Pendulum

The Period of a conical pendulum  $(T) = 2\pi \sqrt{(l \cos \theta / g)}$

where

l is the length of the pendulum

g is acceleration due to gravity

Half angle of the conical pendulum

## Electricity

---

Here are some formulas related to electricity.

## Ohm's Law

---

Ohm's law gives a relation between the voltage applied a current flowing across a solid conductor:

$V$  (Voltage) =  $I$  (Current) x  $R$  (Resistance)

## Power

---

In case of a closed electrical circuit with applied voltage  $V$  and resistance  $R$ , through which current  $I$  is flowing,

Important topics of physics with equations

Power (P) =  $V^2$

R

=  $I^2R$ . . . (because  $V = IR$ , Ohm's Law)

## Kirchoff's Voltage Law

---

For every loop in an electrical circuit:

$$\sum_i V_i = 0$$

where  $V_i$  are all the voltages applied across the circuit.

Kirchoff's Current Law

At every node of an electrical circuit:

$$\sum_i I_i = 0$$

where  $I_i$  are all the currents flowing towards or away from the node in the circuit.

## Resistance

---

The physics formulas for equivalent resistance in case of parallel and series combination are as follows:

Resistances  $R_1$ ,  $R_2$ ,  $R_3$  in series:

Important topics of physics with equations

$$R_{eq} = R_1 + R_2 + R_3$$

Resistances  $R_1$  and  $R_2$  in parallel:

$$R_{eq} = \frac{R_1 R_2}{R_1 + R_2}$$

$$R_1 + R_2$$

For  $n$  number of resistors,  $R_1, R_2, \dots, R_n$ , the formula will be:

$$\frac{1}{R_{eq}} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots + \frac{1}{R_n}$$

## Capacitors

---

A capacitor stores electrical energy, when placed in an electric field. A typical capacitor consists of two conductors separated by a dielectric or insulating material. Here are the most important formulas related to capacitors. Unit of capacitance is Farad (F) and its values are generally specified in mF (micro Farad =  $10^{-6}$  F).

$$\text{Capacitance (C)} = Q / V$$

$$\text{Energy Stored in a Capacitor (E}_{cap}) = \frac{1}{2} CV^2 = \frac{1}{2} (Q^2 / C) = \frac{1}{2} (QV)$$

$$\text{Current Flowing Through a Capacitor } I = C (dV / dt)$$

## Important topics of physics with equations

Equivalent capacitance for 'n' capacitors connected in parallel:

$$C_{eq} (\text{Parallel}) = C_1 + C_2 + C_3 \dots + C_n = \sum_{i=1}^n C_i$$

Equivalent capacitance for 'n' capacitors in series:

$$1 / C_{eq} (\text{Series}) = 1 / C_1 + 1 / C_2 \dots + 1 / C_n = \sum_{i=1}^n (1 / C_i)$$

Here

C is the capacitance

Q is the charge stored on each conductor in the capacitor

V is the potential difference across the capacitor

Parallel Plate Capacitor Formula:

$$C = k\epsilon_0 (A/d)$$

Where

k = dielectric constant (k = 1 in vacuum)

$\epsilon_0$  = Permittivity of Free Space (=  $8.85 \times 10^{-12} \text{ C}^2 / \text{Nm}^2$ )

A = Plate Area (in square meters)

d = Plate Separation (in meters)

Cylindrical Capacitor Formula:

$$C = 2\pi k\epsilon_0 [L / \ln(b / a)]$$

## Important topics of physics with equations

Where

$k$  = dielectric constant ( $k = 1$  in vacuum)

$\epsilon_0$  = Permittivity of Free Space ( $= 8.85 \times 10^{-12} \text{ C}^2 / \text{Nm}^2$ )

$L$  = Capacitor Length

$a$  = Inner conductor radius

$b$  = Outer conductor radius

Spherical Capacitor Formula:

$$C = 4\pi k\epsilon_0 \left[ \frac{ab}{b-a} \right]$$

Where

$k$  = dielectric constant ( $k = 1$  in vacuum)

$\epsilon_0$  = Permittivity of Free Space ( $= 8.85 \times 10^{-12} \text{ C}^2 / \text{Nm}^2$ )

$a$  = Inner conductor radius

$b$  = Outer conductor radius

Inductors

An inductor is an electrical component which resists the flow of electrons or electric current through it. This property of inductance, in these devices, is caused by the electromotive force, created by magnetic field induced in them. The unit of inductance is Henry. Here are some important formulas associated with inductors.

$$\text{Energy Stored in Inductor (E}_{\text{stored}}) = \frac{1}{2} (LI^2)$$

Where,  $L$  is inductance and  $I$  is the current flowing through the inductor.

$$\text{Inductance of a cylindrical air core Coil (L)} = \frac{\mu_0 k N^2 A}{l}$$

## Important topics of physics with equations

Where

L is inductance measured in Henries

N is the number of turns on the coil

A is cross-sectional area of the coil

$\mu_0$  is the permeability of free space ( $= 4\pi \times 10^{-7} \text{ H/m}$ )

K is the Nagaoka coefficient

l is the length of coil

Inductors in a Series Network

For inductors,  $L_1, L_2 \dots L_n$  connected in series,

$$L_{eq} = L_1 + L_2 \dots + L_n \text{ (L is inductance)}$$

Inductors in a Parallel Network

For inductors,  $L_1, L_2 \dots L_n$  connected in parallel,

$$1 / L_{eq} = 1 / L_1 + 1 / L_2 \dots + 1 / L_n$$

# Thermodynamics Formulas

---

## Important topics of physics with equations

Thermodynamics is a vast field providing an analysis of the behavior of matter in bulk. It's a field focused on studying matter and energy in all their manifestations. Here are some of the most important formulas associated with classical thermodynamics and statistical physics.

### First Law of Thermodynamics

$$dU = dQ + dW$$

where,  $dU$  is the change in internal energy,  $dQ$  is the heat absorbed by the system and  $dW$  is the work done on the system.

# Thermodynamic Potentials

---

All of thermodynamical phenomena can be understood in terms of the changes in five thermodynamic potentials under various physical constraints. They are Internal Energy ( $U$ ), Enthalpy ( $H$ ), Helmholtz Free Energy ( $F$ ), Gibbs Free Energy ( $G$ ), Landau or Grand Potential ( $\Phi$ ). Each of these scalar quantities represents the potentiality of a thermodynamic system to do work of various kinds under different types of constraints on its physical parameters.

### Thermodynamic Potential Defining Equation

$U$  is Energy

$T$  is temperature

$S$  is Entropy

$N$  is particle number

$\mu$  is Chemical Potential

$p$  is Pressure

$V$  is Volume

$H$  is Enthalpy

## Important topics of physics with equations

G is Gibbs Free Energy

$\Phi$  is Grand Potential

F is Helmholtz Free Energy

Internal Energy (U)  $dU = TdS - pdV + \mu dN$

Enthalpy (H)  $H = U + pV$

$dH = TdS + Vdp + \mu dN$

Gibbs Free Energy (G)  $G = U - TS + pV = F + pV = H - TS$

$dG = -SdT + Vdp + \mu dN$

Helmholtz Free Energy (F)  $F = U - TS$

$dF = -SdT - pdV + \mu dN$

Landau or Grand Potential  $\Phi = F - \mu N$

$d\Phi = -SdT - pdV - Nd\mu$

# Ideal Gas Equations

---

An ideal gas is a physicist's conception of a perfect gas composed of non-interacting particles which are easier to analyze, compared to real gases, which are much more complex, consisting of interacting particles. The resulting equations and laws of an ideal gas conform with the nature of real gases under certain conditions, though they fail to make exact predictions due to the interactivity of molecules, which is not taken into consideration. Here are some of the most important physics formulas and equations, associated with ideal gases. Let's begin with the prime ideal gas laws and the equation of state of an ideal gas.

## Law Equation

---

P is Pressure

V is Volume

## Important topics of physics with equations

T is Temperature

n is the number of moles

R is the ideal gas constant [= 8.3144621(75) J / K mol]

N is the number of particles

k is the Boltzmann Constant (= 1.3806488(13)×10<sup>-23</sup>)

Boyle's Law  $PV = \text{Constant}$

or

$$P_1V_1 = P_2V_2$$

(At Constant Temperature)

Charles's Law  $V / T = \text{Constant}$

or

$$V_1 / T_1 = V_2 / T_2$$

(At Constant Pressure)

Amontons' Law of Pressure-Temperature  $P / T = \text{Constant}$

or

$$P_1 / T_1 = P_2 / T_2$$

(At Constant Volume)

Equation of State For An Ideal Gas  $PV = nRT = NkT$

# Kinetic Theory of Gases

---

Based on the primary assumptions that the volume of atoms or molecules is negligible, compared to the container volume and the attractive forces between molecules are negligible, the kinetic theory describes the properties of ideal gases. Here are the most important physics formulas related to the kinetic theory of monatomic gases.

## Important topics of physics with equations

$$\text{Pressure (P)} = \frac{1}{3} (Nm v^2)$$

Here, P is pressure, N is the number of molecules and  $v^2$  is the mean squared particle velocity.

$$\text{Internal Energy (U)} = \frac{3}{2} (NkT)$$

### Heat Capacities

$$\text{Heat Capacity at Constant Pressure (Cp)} = \frac{5}{2} Nk = C_v + Nk$$

$$\text{Heat Capacity at Constant Volume (Cv)} = \frac{3}{2} Nk$$

$$\text{Ratio of Heat Capacities } (\gamma) = C_p / C_v = 5/3$$

### Velocity Formulas

$$\text{Mean Molecular Velocity (Vmean)} = [(8kT)/(\pi m)]^{1/2}$$

$$\text{Root Mean Square Velocity of a Molecule (Vrms)} = (3kT/m)^{1/2}$$

$$\text{Most Probable Velocity of a Molecule (Vprob)} = (2kT/m)^{1/2}$$

$$\text{Mean Free Path of a Molecule } (\lambda) = (kT)/\sqrt{2}nd^2P \text{ (Here P is in Pascals)}$$

Important topics of physics with equations

Here  $N$  is the number of molecules,  $k$  is the Boltzmann constant,  $P$  is pressure,  $d$  is the molecular diameter,  $m$  is mass of the molecule and  $T$  is the gas temperature.

# Electromagnetism

---

Here are some of the basic formulas from electromagnetism.

The coulombic force between two charges at rest is

$$F = \frac{q_1 q_2}{4\pi\epsilon_0 r^2}$$

$$4\pi\epsilon_0 r^2$$

Here,

$q_1, q_2$  are charges

$\epsilon_0$  is the permittivity of free space

$r$  is the distance between the two charges

Lorentz Force

The Lorentz force is the force exerted by an electric and/or magnetic field on a charged particle.

$$\text{(Lorentz Force) } F = q (E + v \times B)$$

where

$q$  is the charge on the particle

$E$  and  $B$  are the electric and magnetic field vectors

Relativistic Mechanics

Here are some of the most important relativistic mechanics formulas. The transition from classical to relativistic mechanics is not at all smooth, as it merges space and time into one by taking away the Newtonian idea of absolute time. If you know what is Einstein's special theory of relativity, then the following formulas will make sense to you.

# Lorentz Transformations

---

Lorentz transformations can be perceived as rotations in four dimensional space. Just as rotations in 3D space mixes the space coordinates, a Lorentz transformation mixes time and space coordinates. Consider two, three dimensional frames of reference  $S(x,y,z)$  and  $S'(x',y',z')$  coinciding with each other.

Now consider that frame  $S'$  starts moving with a constant velocity  $v$  with respect to  $S$  frame. In relativistic mechanics, time is relative! So the time coordinate for the  $S'$  frame will be  $t'$  while that for  $S$  frame will be  $t$ .

Consider

$$\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$$

$$\sqrt{1 - v^2/c^2}$$

The coordinate transformations between the two frames are known as Lorentz transformations and are given as follows:

Lorentz Transformations of Space and Time

$$x = \gamma (x' + vt')$$
$$x' = \gamma (x - vt)$$

$$y = y'$$

$$z = z'$$

$$t = \gamma (t' + vx'/c^2)$$
$$t' = \gamma (t - vx/c^2)$$

### Relativistic Velocity Transformations

In the same two frames S and S', the transformations for velocity components will be as follows (Here  $(U_x, U_y, U_z)$  and  $(U_x', U_y', U_z')$  are the velocity components in S and S' frames respectively):

$$U_x = (U_x' + v) / (1 + U_x'v / c^2)$$

$$U_y = (U_y') / \gamma(1 + U_x'v / c^2)$$

$$U_z = (U_z') / \gamma(1 + U_x'v / c^2) \text{ and}$$

$$U_x' = (U_x - v) / (1 - U_xv / c^2)$$

$$U_y' = (U_y) / \gamma(1 - U_xv / c^2)$$

$$U_z' = (U_z) / \gamma(1 - U_xv / c^2)$$

## Momentum and Energy Transformations in Relativistic Mechanics

---

Consider the same two frames (S, S') as in case of Lorentz coordinate transformations above. S' is moving at a velocity 'v' along the x-axis. Here again  $\gamma$  is the Lorentz factor. In S frame  $(P_x, P_y, P_z)$  and in S' frame  $(P_x', P_y', P_z')$  are momentum components. Now we consider formulas for momentum and energy transformations for a particle, between these two reference frames in relativistic regime.

## Important topics of physics with equations

### Component wise Momentum Transformations and Energy Transformations

$$P_x = \gamma(P_x' + vE' / c^2)$$

$$P_y = P_y'$$

$$P_z = P_z'$$

$$E = \gamma(E' + vP_x')$$

and

$$P_x' = \gamma(P_x - vE / c^2)$$

$$P_y' = P_y$$

$$P_z' = P_z$$

$$E' = \gamma(E - vP_x)$$

### Physical Formulas for Quantities in Relativistic Dynamics

All the known quantities in classical mechanics get modified, when we switch over to relativistic mechanics which is based on the special theory of relativity. Here are formulas of quantities in relativistic dynamics.

Important topics of physics with equations

Relativistic momentum  $p = \gamma m_0 v$

where  $m_0$  is the rest mass of the particle.

Rest mass energy  $E = m_0 c^2$

Total Energy (Relativistic)  $E = \sqrt{(p^2 c^2 + m_0^2 c^4)}$

## Optics

---

Optics is one of the oldest branches of physics. There are many important optics physics formulas, which we need frequently in solving physics problems. Here are some of the important and frequently needed optics formulas.

## Snell's Law

---

$\sin i$

$\sin r = n_2$

$n_1 = v_1$

$v_2$

where  $i$  is angle of incidence

$r$  is the angle of refraction

$n_1$  is refractive index of medium 1

$n_2$  is refractive index of medium 2

## Important topics of physics with equations

$v_1, v_2$  are the velocities of light in medium 1 and medium 2 respectively

Gauss Lens Formula:  $1/u + 1/v = 1/f$

where

$u$  - object distance

$v$  - image distance

$f$  - Focal length of the lens

Lens Maker's Equation

The most fundamental property of any optical lens is its ability to converge or diverge rays of light, which is measured by its focal length. Here is the lens maker's formula, which can help you calculate the focal length of a lens, from its physical parameters.

$$1/f = [n-1][(1/R_1) - (1/R_2) + (n-1)d/nR_1R_2]$$

Here,

$n$  is refractive index of the lens material

$R_1$  is the radius of curvature of the lens surface, facing the light source

$R_2$  is the radius of curvature of the lens surface, facing away from the light source

$d$  is the lens thickness

If the lens is very thin, compared to the distances -  $R_1$  and  $R_2$ , the above formula can be approximated to:

$$\text{(Thin Lens Approximation) } 1/f \approx (n-1) [1/R_1 - 1/R_2]$$

Compound Lenses

## Important topics of physics with equations

The combined focal length ( $f$ ) of two thin lenses, with focal length  $f_1$  and  $f_2$ , in contact with each other:

$$1 / f = 1 / f_1 + 1 / f_2$$

If the two thin lenses are separated by distance  $d$ , their combined focal length is provided by the formula:

$$1 / f = 1 / f_1 + 1 / f_2 - (d / f_1 f_2)$$

### Newton's Rings Formulas

Here are the important formulas for Newton's rings experiment which illustrates diffraction.

$n$ th Dark ring formula:  $r^2_n = nR\lambda$

$n$ th Bright ring formula:  $r^2_n = (n + \frac{1}{2}) R\lambda$

where

$n$ th ring radius

Radius of curvature of the lens

Wavelength of incident light wave

[[Back to Formula Index](#)]

Important topics of physics with equations

Quantum Physics

Quantum physics is one of the most interesting branches of physics, which describes atoms and molecules, as well as atomic sub-structure. Here are some of the formulas related to the very basics of quantum physics, that you may require frequently.

## De Broglie Wave

---

De Broglie Wavelength:

$$\lambda = \frac{h}{p}$$

p

where,  $\lambda$ - De Broglie Wavelength, h - Planck's Constant, p is momentum of the particle.

Bragg's Law of Diffraction:  $2a \sin \theta = n\lambda$

where

a - Distance between atomic planes

n - Order of Diffraction

$\theta$  - Angle of Diffraction

$\lambda$  - Wavelength of incident radiation

Planck Relation

The plank relation gives the connection between energy and frequency of an electromagnetic wave:

$$E = h\nu = h\omega$$

$2\pi$

Important topics of physics with equations

where  $h$  is Planck's Constant,  $\nu$  the frequency of radiation and  $\omega = 2\pi\nu$

## Uncertainty Principle

---

Uncertainty principle is the bedrock on which quantum mechanics is based. It exposes the inherent limitation that nature imposes on how precisely a physical quantity can be measured. Uncertainty relation holds between any two non-commuting variables. Two of the special uncertainty relations are given below.

## Position-Momentum Uncertainty

---

What the position-momentum uncertainty relation says is, you cannot predict where a particle is and how fast it is moving, both, with arbitrary accuracy. The more precise you are about the position, more uncertain will you be about the particle's momentum and vice versa. The mathematical statement of this relation is given as follows:

$$\Delta x \cdot \Delta p \geq \frac{h}{2\pi}$$

$$2\pi$$

where  $\Delta x$  is the uncertainty in position and  $\Delta p$  is the uncertainty in momentum.

## Energy-Time Uncertainty

---

This is an uncertainty relation between energy and time. This relation gives rise to some astounding results like, creation of virtual particles for arbitrarily short periods of time! It is mathematically stated as follows:

$$\Delta E \cdot \Delta t \geq \frac{h}{2\pi}$$

$$2\pi$$

## Important topics of physics with equations

where  $\Delta E$  is the uncertainty in energy and  $\Delta t$  is the uncertainty in time.

This concludes my review of some of the important physics formulas. This list, is only representative and is by no means anywhere near complete. Physics is the basis of all sciences and therefore its domain extends over all sciences. Every branch of physics theory abounds with countless formulas. If you resort to just mugging up all these formulas, you may pass exams, but you will not be doing real physics. If you grasp the underlying theory behind these formulas, physics will be simplified. To view physics through the formulas and laws, you must be good at maths. There is no way you can run away from it. Mathematics is the language of nature!

The more things we find out about nature, more words we need to describe them. This has led to increasing jargonization of science with fields and sub-fields getting generated. You could refer to a glossary of science terms and scientific definitions for any jargon that is beyond your comprehension.

If you really want to get a hang of what it means to be a physicist and get an insight into physicist's view of things, read 'Feynman Lectures on Physics', which is highly recommended reading, for anyone who loves physics. It is written by one of the greatest physicists ever, Prof. Richard Feynman. Read and learn from the master. Solve as many problems as you can, on your own, to get a firm grasp of the subject.